



Melatonin; an Emerging Physiological Regulator in Postharvest Management of Horticultural Crops: Mechanisms and Practical Applications

Mani Jabbari^{1*}, Mitra Jabbari²

¹ Department of Horticultural Science and Engineering, Faculty of Agriculture, Birjand University, Iran.

² Department of Horticultural Science and Engineering, Faculty of Plant Production, Gorgan University of Agricultural Sciences and Natural Resources

Abstract

The increase in postharvest losses in horticultural crops, especially fruits and vegetables, has become one of the major challenges in food security and supply chain sustainability due to high perishability, physiological stresses, and oxidative damage. The use of traditional chemical compounds, although effective to some extent in spoilage control, has highlighted the need for safe and biocompatible alternatives due to environmental concerns and consumer health risks. In the meantime, melatonin (N-acetyl-5-methoxytryptamine) as an endogenous, non-toxic, and multifunctional signaling molecule has great potential for application in postharvest technology. Research evidence shows that exogenous application of melatonin can prevent the occurrence of physiological disorders and premature aging by strengthening the antioxidant defense system, increasing the activity of protective enzymes and non-enzymatic compounds, reducing the accumulation of reactive oxygen species, and maintaining the integrity of cell membranes. In addition, melatonin modulates the expression of defense genes and increases the tolerance of horticultural products to post-harvest stresses, including frost, by regulating ethylene-, salicylic acid-, and abscisic acid-dependent signaling pathways. These properties maintain quality traits such as tissue firmness, soluble matter content, phenolic compounds, and vitamins during storage. Accordingly, melatonin can be a suitable alternative to conventional chemical methods as a green and sustainable preservative and play an effective role in reducing waste, increasing shelf life, and improving the quality of horticultural products.

Keywords: Antioxidant, Aging, Frost Damage, Photosynthesis, Phenolic, Fruit Quality.

* Received 29 November 2025; Received in revised from 25 December 2025; Accepted 28 October 2025; Published online Jan 2026.

Cite this article: Jabbari, M & Jabbari, M. (2026). Melatonin; an emerging physiological regulator in postharvest management of horticultural crops: mechanisms and practical applications, *Iranian journal of postharvest biology*, 2(4), 102-123. [HTTPS://DOI.ORG/10.22091/IJPB.2026.16926.1014](https://doi.org/10.22091/IJPB.2026.16926.1014)

Publisher: University of Qom



© The Author(s)

1. Introduction

The growing global population has significantly increased the demand for food (Ema et al., 2024). A certain percentage of food produced globally is lost or wasted. (Lau et al., 2021), is a serious challenge in the food supply chain. Several post-harvest technologies, including cold storage and the use of synthetic chemicals, have been applied to reduce this waste and improve product quality (Ferreira et al., 2024; Patil et al., 2024; Zhang, 2024). However, due to increasing environmental and health concerns, the use of synthetic chemicals is decreasing (Bose et al., 2021), which highlights the need to develop new and sustainable technologies. In this regard, exogenous plant hormone treatments have been proposed as a natural and promising solution to maintain the quality and increase the shelf life of fruits and vegetables (Aghdam et al., 2020; Baswal et al., 2020; Adhikary et al., 2021; Islam et al., 2022; Gutiérrez-Villamil et al., 2023). Phytohormones, natural organic compounds, play a pivotal role in coordinating plant physiological activities, responding to environmental stresses, and regulating growth and development (Hu et al., 2022; Massolo et al., 2022). Horticultural crops, due to their vital role in food security, commercial economy, and industrial applications, have a special place in global agriculture and are considered a rich source of micronutrients, vitamins, and bioactive compounds (Gao et al., 2022). However, the production of these products is always constrained by biotic (e.g., pests and pathogens) and abiotic (e.g., temperature fluctuations, drought, salinity, and heavy metals) stresses, which lead to a decrease in the quantity and quality of products (Ahmad et al., 2019; Kumari et al., 2021; Umamaheswari et al., 2020). Therefore, it is essential to develop effective management strategies to increase the resilience and sustainability of the production of these products. The high perishability of horticultural products due to mechanical damage, physiological deterioration, and improper post-harvest management is a serious obstacle in their supply chain and marketing (Xu et al., 2019). Despite the widespread use of technologies and chemicals to control spoilage, their environmental and health-

related risks have drawn researchers' attention to natural preservatives (Bose et al., 2021). In response to the increasing demand for safe and nutritious products, the use of compounds such as melatonin has attracted special attention as a novel and sustainable solution to increase the shelf life and maintain the quality of horticultural products (Nitu et al., 2024; Padilla-González et al., 2026).

Melatonin (N-acetyl-5-methoxytryptamine, MT), an indole compound first identified in the pineal gland of animals (Wang et al., 2026; Favero et al., 2026; Huang and Wong, 2026), has been widely considered in recent years as an effective tool in promoting stress tolerance and improving plant growth (Ahmad et al., 2023). This compound, which is endogenously present in many plant species, plays a key role in regulating physiological activities (Arnao and Hernández-Ruiz, 2014). Research has shown that exogenous application of melatonin, in addition to stimulating growth, increases tolerance to environmental stresses and delays the aging process by modulating defense responses (Zhou et al., 2021; Wang et al., 2023; Nguyen and Ha, 2025).




Figure 1.
Chemical structure of Melatonin (MEL). (Padilla-González et al., 2026).

Melatonin, as a plant growth regulator (PGR), affects important processes such as fruit formation, seed germination, root proliferation, flowering, and fruit ripening (Dai et al., 2020). In plants, melatonin acts as a biological signal in several plant processes such as photosynthesis, growth, defense, germination, budding, flowering, leaf senescence, membrane integrity, root growth, osmotic regulation, and plant protection against biotic and abiotic stresses (Sharma and Zheng, 2019). Also, this compound is able to increase the shelf life and freshness of fruits after

harvest. The role of melatonin in stress resistance, hormone regulation, and postharvest quality improvement has been consistently reported (Wang et al., 2020; Jabbari and Jabbari, 2023). Treatment with exogenous melatonin can improve the stress resistance, antioxidant capacity, and antimicrobial ability of fruits after harvest, thereby increasing their shelf life and quality (Hernández-Ruiz et al., 2022). In particular, the function of melatonin in delaying fruit ripening and senescence during storage has attracted the attention of researchers, as the mechanism of its influence on the quality of harvested fruits has become an important research focus (Arnao and Hernández-Ruiz, 2020; Kumar Bose and Howlade, 2026). Melatonin has recently been studied to maintain postharvest quality and increase the shelf life of various horticultural crops (Hei et al., 2025). External application of melatonin can enhance endogenous melatonin to enhance its activities in several physiological processes and scavenge Reactive Oxygen Species (ROS) from fruits and vegetables during storage by increasing the activity of antioxidant enzymes. Several previous reports have indicated that external application of melatonin can maintain quality, control postharvest diseases, and extend the shelf life of jujube (Hei et al., 2025), plum (Zhang et al., 2024), pepper (Li et al., 2024; Charoenphun et al., 2024), guava (Yang et al., 2023), papaya (Wang et al., 2022), sweet cherry (Wang et al., 2019; Bal, 2023), strawberry (Liu et al., 2018), tomato (Sharafi et al., 2019), grape (Meng et al., 2019), peach (Gao et al., 2018), pear (Liu et al., 2019) and many others have gradually attracted the attention of researchers and consumers. In this review, we systematically summarize recent research advances that reveal the multiple functions and mechanisms of action of melatonin during postharvest storage of horticultural crops and provide future research directions for the use of melatonin for postharvest management.

2. Melatonin metabolism in plants

Melatonin (N-acetyl-5-methoxytryptamine) is an indoleamine compound derived from the amino acid tryptophan that is found in a wide range of organisms, from microorganisms to vertebrates (Juhnevica-Radenkova et al., 2020). It was first

isolated and identified in 1958 from the bovine pineal gland.

In the early decades after its discovery, melatonin was known solely as a neurohormone in mammals, and studies focused on its role in regulating circadian rhythms, modulating immune responses, and controlling seasonal reproductive cycles; however, with the development of biochemistry, the functional scope of this hormone is now being evaluated far beyond these classical definitions (Shao et al., 2024).

As research progressed, melatonin was also found in other biological domains. In the 1980s, it was reported in invertebrates, and in 1989, it was first identified in a phototrophic unicellular organism called *Lingulodinium polyedrum* (formerly *Gonyaulax polyedra*). Finally, findings in 1995 showed that melatonin is also present in plant species, thus introducing this compound as a biological molecule with a wide distribution in nature (Salehi et al., 2019). In plants, melatonin is a multifunctional compound whose biosynthesis occurs through the Shikimate metabolic pathway using the amino acid tryptophan as the main precursor. This biosynthetic process is complex and takes place in various cellular organelles, including the cytoplasm, chloroplasts, mitochondria, and endoplasmic reticulum (Kumari et al., 2023).




Figure 2.

Melatonin (MT) acts as a dual-organelle antioxidant shield against ROS (Eisa et al., 2026).

The main pathway of melatonin biosynthesis in plants involves four sequential enzymatic reactions: tryptophan is decarboxylated by the enzyme tryptophan decarboxylase (TDC) and converted to tryptamine; tryptamine is converted to serotonin (5-hydroxytryptamine) by the enzyme tryptamine-5-

hydroxylase (T5H); serotonin is converted to N-acetyl serotonin (NAS) by the enzyme serotonin-N-acetyltransferase (SNAT); in the final step, NAS is methylated by the activity of the enzymes N-acetyl serotonin methyltransferase (ASMT) or caffeic acid-O-methyltransferase (COMT) and converted to melatonin (He et al., 2023; Chen et al., 2026). The melatonin biosynthesis pathway in plants is linked to the synthesis of other secondary metabolites and is influenced by a complex network of genetic factors, plant hormones, and environmental stresses.

Previously, several researchers have identified melatonin biosynthesis genes in various horticultural crops such as apple, cherry, pepper, tomato, and watermelon, suggesting that melatonin is endogenously biosynthesized and plays an important role in growth and defense responses. For example, the MdTDC, MdT5H, MdAANAT, and MdASMT genes were identified in the fruit of 'Red Fuji' apple and are involved in melatonin biosynthesis (Lei et al., 2013). In another study, the PaTDC gene was identified in cherry fruit, and it was suggested that PaTDC positively regulates melatonin biosynthesis (Yu Zhao et al., 2012). SlCOMT1 is another gene identified in tomato fruit (*Solanum lycopersicum* Mill. cv. Ailsa Craig) (Liu et al., 2019). Various genes such as ClTDC, ClT5H, ClSNAT, ClASMT, and ClCOMT are involved in melatonin biosynthesis in watermelon (Mandal et al., 2018). Therefore, the above findings indicate that melatonin is endogenously produced in different parts of plants including fruits, flowers, leaves, seeds, etc.




Figure 3.

Melatonin biosynthesis pathways in horticulture crops (Gao et al., 2022).

Table 1.
Effect of melatonin on different fruit crops

Scientific name	Application method	Effective concentration ($\mu\text{M/ppm}$)	Plant response	References
<i>Mangifera indica</i> L.	Immersed	500 μM	Delayed ripening and softening	Liu et al., 2020
<i>Musa acuminata</i> L.	Immersed	200 or 500 μM	Delayed postharvest ripening	Hu et al., 2017
Kiwifruit	Immersed	50 μM	Enhanced their germination rate and germination potential	Shen et al., 2019
Apple	Sprayed	–	Promoted adventitious root formation mainly at the stage of AR induction by increasing IAA levels and activating the function of MdWOX11 (overexpression lines)	Mao et al., 2020
Grapes (5BB rootstock and the cultivar Cabernet Sauvignon)	Immersed	0.1 and 0.5 μM	Induced rooting in cultivar and root-stock	Gökbayrak et al., 2020
Pomegranate	sprayed	5 ppm	Reduces the fruit drop and viable hermaphrodite flowers drop, which increases the fruit set	Usanmaz et al., 2022
Apple	sprayed	20 and 200 μM	Improved flowering rate	Zhang et al., 2019
Apricot	sprayed	25 ppm	Increased vegetative growth	El-Naby et al., 2019
Sweet cherry	sprayed	0.3 mM	Improved crop yield and quality	Carrión-Antolí et al., 2021
Pomegranate	sprayed	0.1 mM	Increased crop yield as well as fruit quality	Medina-Santamarina et al., 2021
Grape (Kyoho)	Immersed	200 μM	Reduced berry abscission and rotten index	Wang et al., 2019
Grape (Moldova)	Immersed	10 and 100 μM	Promoted berry ripening	Xu et al., 2018
Banana	Immersed	–	Delayed chilling injury and alleviating peel browning	Cao et al., 2018
Jujube	sprayed	200 μmolL^{-1}	Increased the disease resistance against <i>Alternaria rot</i>	Zhang et al., 2022
Apple (golden delicious)	Immersed	50 μmolL^{-1}	Induced resistance to blue mold	Wang et al., 2023

Table 2.
Melatonin concentration in different fruit species

Fruits	Scientific name	Variety/cultivar	Concentration (ng g ⁻¹)	Stage of evaluation	References
Almond	<i>Prunus dulcis</i>	–	0.12	Mature	Paroni et al., 2019
Red grape (wine)	<i>Vitis vinifera</i> L.	Nebbiolo	0.57 to 0.63	Ripe	Fracassetti et al., 2019
		“Nero d’Avola	0.05 to 0.62	Ripe	Fracassetti et al., 2019
Sour cherry	<i>Prunus ceracus</i>	–	20	In the Second stage of fruit development	Tijero et al., 2019
Goji berry/wolfberry	<i>Lycium barbarum</i>		1600.48 ± 23.66	Mature	Yılmaz, 2021
Black mulberry	<i>Morus nigra</i>		178.70 ± 0.57		
White mulberry	<i>Morus alba</i>		183.29 ± 6.27		
Blackberry	<i>Rubus fruticosus</i> L.		177.91 ± 27.51		
Purple mulberry	<i>Morus rubra</i>		233.86 ± 7.20		
Bursa black mulberry	–		123.44 ± 2.05		
Cornelian cherry	<i>Cornus mas</i> L.	–	130.82—201.84	Mature	Uğur, 2023
Date Palm	<i>Phoenix dactylifera</i>	Deglet Nour	0.02	Mature	Verde et al., 2019
		Allighes	0.0074		
		Medjbouda	0.02		
		Kenta	0.17		
		Khadrawy	0.0148		
		Medjou	0.0087		
Walnut	<i>Juglans regia</i>	Hartley	3.3	Mature	Verde et al., 2022
		Franquette	1.6		
		Pizarro	1.19		
		Native	2.07		
Apple	<i>Malus domestica</i> Mill.	Granny Smith	7.37	Ripe	Zhang et al., 2018
		Fuji	67.63		
Mulberry	<i>Morus alba</i>	–	1.739 ± 0.076	Mature	Pranil et al., 2021
Guava	<i>Psidium guajava</i>	–	1.570 ± 0.015	Mature	

Banana	<i>Musa sapientum</i>	—	1.741 ± 0.103	Ripe
Mango	<i>Mangifera indica</i>	—	2.401 ± 0.036	Mature
Pineapple	<i>Ananas comosus</i>	—	1.693 ± 0.088	Mature
Orange	<i>Citrus sinensis</i>	—	1.704 ± 0.028	Mature
Longan	<i>Dimocarpus longan</i>	—	1.676 ± 0.080	Mature
Red passion fruit	<i>Passiflora edulis</i>	—	1.528 ± 0.113	Mature
Yellow passion fruit	<i>Passiflora edulis</i>	—	1.744 ± 0.140	Mature
Jujube	<i>Ziziphus jujuba</i>	—	1.679 ± 0.163	Mature

3. Increasing shelf life and improving post-harvest quality

The use of melatonin as a post-harvest treatment plays an effective role in maintaining the quality and increasing the shelf life of agricultural products. In a study on nectarine (Fantasia cultivar), immersing fruits in different concentrations of melatonin (250, 500, and 1000 μmolL^{-1}) and storing at temperatures of 0 to 1 °C and relative humidity of 85 to 90% resulted in maintaining the texture (firmness) of the fruit and preventing browning of its skin and flesh. In addition, this treatment, while reducing respiration rate and weight loss, favorably maintained quality indicators such as total soluble solids (TSS) and soluble sugars (Chen et al., 2022). In another study on cassava root tubers, exogenous melatonin significantly prevented postharvest physiological deterioration (PPD) by modulating oxidative stress through reducing hydrogen peroxide levels and increasing the activities of antioxidant enzymes catalase and peroxidase (Hu et al., 2016). Overall, the available evidence suggests that melatonin application is an effective strategy to delay the aging process and reduce postharvest losses in horticultural crops, which is achieved through improving antioxidant defense mechanisms and maintaining physicochemical properties.

4. Effect of melatonin on the quality of harvested fruits

The visual quality of fruits includes factors such as size, firmness, skin smoothness, color, and other aspects. These visual characteristics directly affect the commercial value of the fruit and also influence consumer choice (Li et al., 2023; Shi et al., 2024). Studies have shown that melatonin enhances antioxidant systems and improves membrane integrity, thereby reducing chilling injury and increasing fruit firmness and color (Padilla-González et al., 2026). Melatonin (100 μmolL^{-1}) delayed the loss of firmness of guava fruit and reduced fruit weight loss (Chen et al., 2022). Spraying melatonin at a concentration of 100 μmolL^{-1} on grapes during the variation period induced skin pigmentation earlier than the control group and also caused higher transcript abundance of genes related to anthocyanin

biosynthesis and the transcription factors MYBA1 and MYBA2 (Xia et al., 2021).

5. Slowing down ripening and delaying senescence

Melatonin treatments delay ripening by reducing softening, weight loss, ethylene production and respiration rate (Zhai et al., 2018; Rastegar et al., 2020; Onik et al., 2021). This leads to increased TSS, TA and inhibition of surface browning (Aghdam and Fard, 2017; Xu et al., 2018; Liu et al., 2018; Zheng et al., 2019; Fan et al., 2023). The efficacy of melatonin is strongly linked to its ability to suppress ROS production by increasing the activity of antioxidant enzymes (Zhang et al., 2018; Ze et al., 2021). For example, Gao et al. (2016) reported increased APX, SOD, POD, and CAT activities in peach fruit treated with 100 μM melatonin during 7 days of storage at ambient temperature. Similar results were observed in pomegranate fruit stored at 4°C for 120 days with the same melatonin concentration (Seyed et al., 2020). Melatonin treatments are significantly associated with increases in nonenzymatic antioxidant systems, including anthocyanins, flavonoids, and ascorbic acid (Sun et al., 2020; Seyed et al., 2020).

Table 3.

Several studies on the effect of exogenous melatonin on the postharvest treatment of fruits

Botanical Name	Optimum Concentration (μM)	Effects	Application	References
<i>Mangifera indica</i> L.	100	Soluble solids, pH, total soluble solid and acidity ratio; maintained a higher concentration of total phenolics and total flavonoids; increased the activities of antioxidant enzymes superoxide dismutase and CAT; and improved membrane stability.	Dipped for 120 min	Bhardwaj et al., 2022
<i>Mangifera indica</i>	1000	Improved postharvest quality; maintained firmness; improved bioactive compounds; increased phenolic content; and increased the activity of CAT and peroxidase (POD).	Immersed for 10 min	Rastegar et al., 2020
<i>Carica papaya</i>	400	Delayed fruit softening; reduced anthracnose incidence; enhanced antioxidant activity; and decreased fruit oxidative injury; induced total phenol, total flavonoid, and ascorbic acid accumulation; increased CAT, ascorbate peroxidase, NADH oxidase, glutathione reductase, polyphenol oxidase, superoxide dismutase, and peroxidase activity; enhanced the activity of defense-related enzymes, such as chitinase, 4-coumaric acid-CoA-ligase, and phenylalanine ammonia lyase; repressed lipid metabolism.	Immersed for 2 h	Fan et al., 2022
<i>Momordica charantia</i> L.	120	Increased chlorophyll, total soluble solids (TSS), soluble sugar, soluble protein, and ascorbic acid (AsA); promoted the synthesis of total phenols and flavonoids.	Immersed for 10 min	Lin et al., 2020
<i>Vaccinium spp.</i>	50	Maintained the contents of AsA, anthocyanin, and total phenol; reduced the accumulation of reactive oxygen species and membrane lipid peroxidation; and promoted the activities of ascorbate peroxidase (APX), glutathione S-transferase, and phenylalanine ammonia lyase (PAL).	Sprayed	Shang et al., 2021
<i>Musa paradisiaca</i> L.	200–500	Inhibited the ripening process.	Soaked for 2 h	Hu et al., 2017
<i>Psidium guajava</i> L.	600	Maintained postharvest quality, delayed fruit softening, and reduced the incidence of anthracnose; enhanced the antioxidant capacity and reduced the oxidative damage; enhanced the activities of CAT,	Soaked for 2 h	Fan et al., 2022

		superoxide dismutase, ascorbate peroxidase, and glutathione reductase; and maintained total flavonoids and AsA.		
<i>Malus domestica</i> Borkh	1000	Reduced ethylene production and weight loss; maintained skin structure; increased the activity of three enzymes, including POD, superoxide dismutase (SOD), and CAT.	Sprayed	Onik et al., 2021
<i>Malus domestica</i> Borkh	200	Inhibited gray mold diseases.	Immersed	Cao et al., 2017
<i>Prunus persica</i>	100	Reduced weight loss, decay incidence, and respiration rate; maintained firmness, TSS, and AsA contents; enhanced the activities of superoxide dismutase, CAT, peroxidase, and ascorbate peroxidase; decreased the activity of lipoxygenase, levels of superoxide anion and hydrogen peroxide, and malondialdehyde content.	Dipped for 10 min	Gao et al., 2016
<i>Prunus persica</i>	100	Reduced chilling injury; increased extractable juice rate and TSS; enhanced expression of PpADC, PpODC, and PpGAD; increased polyamines and γ -aminobutyric acid (GABA) contents.	Immersed for 2 h	Cao et al., 2016

Other nonenzymatic systems induced by melatonin treatments include carotenoids, dehydroascorbic acid, and glutathione, which contribute to ROS homeostasis (Miranda et al., 2020). Melatonin treatments also induce proline synthesis and increase endogenous melatonin content (Liu et al., 2018; Sharafi et al., 2021;). Notably, melatonin treatments reduce MDA content and EL and LOX enzyme activities while maintaining a higher ratio of unsaturated to saturated fatty acids (Gao et al., 2018; Jannatizadeh et al., 2019). This efficacy is also related to the ability of melatonin to reduce oxidative stress by enhancing antioxidant defense systems.

6. Post-harvest quality and shelf life

Given that the quality of horticultural products after harvest can only be maintained and not improved (Arah et al., 2015), the development of

post-harvest technologies is essential to reduce waste (Ullah et al., 2025). In the meantime, melatonin plays a pivotal role in maintaining the quality of products due to its safe nature and high efficiency (Badiche-El Hilali et al., 2024). Studies show that exogenous melatonin treatment can ensure the nutritional value of products such as jujube and Chinese cabbage by preserving their antioxidant compounds (such as TPC (Total Phenolic Content), TFC (Total Flavonoid Content) and acetylsalicylate (ASA)) and total soluble solids (TSS) (Gurjar et al., 2024; Kang et al., 2025). Furthermore, melatonin significantly increases the physical and chemical shelf life of fruits such as plums, cherries, and papayas by delaying weight loss, softening tissue, and reducing respiration and ethylene production (Zhang et al., 2024; Bal, 2023; Borthakur et al., 2024). These findings make melatonin an ideal option for post-harvest management and reducing horticultural crop waste.

Table 4.

Multifunctional role of melatonin during postharvest storage of horticultural crops

Botanical Name	Application Method	Effective Concentration	Function and Mechanism	References
<i>Anthurium andreaeanum</i>	Dipped	100 μ M	Increased higher phenylalanine ammonia lyase activity and DPPH scavenging activity, ameliorate chilling injury during low temperature storage	Aghdam et al., 2019
<i>Malus domestica</i>	Sprayed	500 μ M	Stimulated fruit ripening through the induction of ethylene synthesis	Verde et al., 2022
<i>Persea americana</i>	Immersed	50 mM	Enhanced total phenol and flavonoid accumulation and maintained titratable acidity, total soluble solids, lipid and ascorbic acid contents	Tu et al., 2022
<i>Musa paradisiaca</i>	Immersed	1000 μ M	Maintained quality and enhanced shelf life during ambient and cold storage	Devi et al., 2025
<i>Vaccinium</i> spp	Soaked	300 μ M	Inhibiting cell wall degradation, improved fruit quality and increased antioxidant capacity	Qu et al., 2022
<i>Brassica oleracea</i> var. italica	Dipped	100 μ M	Maintained higher chlorophyll content and phenolic content during storage	Abou-Elwafa and Safaa, 2025
<i>Manihot esculenta</i>	Soaked	500 mg/L	Delayed postharvest physiological and root deterioration	Hu et al., 2018
<i>Solanum lycopersicum</i> L.	Dipped	0,1 mM	Induced phenylpropanoid pathway and resist against Botrytis cinerea during postharvest storage	Li et al., 2019
<i>Citrus reticulata</i> Blanco	Immersed	50 μ M	Decreases resistance to green mold on citrus fruit by scavenging defense related reactive oxygen species	Lin et al., 2019
<i>Cucumis sativus</i> L.	Immersed	500 μ M	Delayed the loss of chlorophyll, vitamin C, and the content of titratable acidity and soluble protein	Xin et al., 2017
<i>Vitis vinifera</i> L.	Soaked	10 and 100 μ M	Increased the levels of ABA, H ₂ O ₂ , and ethylene production and promoted berry ripening	Xu et al., 2018
<i>Crataegus</i> spp	Immersed	0.10 and 0.25, mM	Maintained fruit quality by increasing phenolic compound during cold storage	Küçüker et al., 2024
<i>Actinidia deliciosa</i>	Immersed	100 and 200 μ M	Inhibited cell wall degradation and preserved fruit quality	Cao et al., 2022
<i>Litchi chinensis</i>	Immersed	0.2 and 0.6	Preserved fruit quality and delayed senescence	Xie et al., 2022

		mM		
<i>Mangifera indica</i>	Immersed	0.5 mM	Reduced chilling injury and enhanced shelf life	Ping et al., 2024
<i>Abelmoschus</i>	Immersed	100 μ M	Delayed senescence by increasing endogenous IAA, GABA, GA and ABA	Shi et al., 2024
<i>Passiflora edulis</i>	Soaked	200, μ M	defense-related mechanisms. Enhanced antioxidant activity and delayed senescence	Cai et al., 2024
<i>Prunus persica</i>	Sprayed	150 μ M	Promotes the sugar accumulation and organic acid degradation in early-ripening	Zhou et al., 2023
<i>Pyrus communis</i>	Immersed	100 μ M	Inhibited ethylene synthesis via nitric oxide regulation and delayed senescence	Jia Zhiwei et al., 2019
<i>Capsicum</i>	Sprayed	100, 200 and 300 μ M	Improved chilling tolerance and enhanced shelf life	Huang et al., 2024
<i>Punica granatum</i>	Immersed	100 μ M	Enhanced antioxidant enzymes activity and maintained nutraceutical properties	Aghdam et al., 2020
<i>Rubus idaeus</i>	Immersed	0.001 and 0.01 mM	Improved phenolic compounds, as well as antioxidant and enzyme activities	Rahmanzadeh-Ishke et al., 2024
<i>Rosa hybrida</i>	Sprayed	20 μ M	Reduced weight loss and bacterial growth	Tongtong et al., 2018
<i>Fragaria ananassa</i>	Immersed	200 and 500 μ M	Maintained postharvest quality during storage	Kahramanog˘lu, 2024
<i>Prunus avium</i>	Immersed	100 and 200 μ M	Delayed softening by reducing cell wall disruption, and maintain the postharvest quality	Ghorbani and Najafzadeh, 2022
<i>Solanum lycopersicum</i> L.	Dipped	100 μ M	Increasing chilling tolerance by enhancing linoleic and linolenic acids accumulation	Jannatizadeh et al., 2019

7. Control of Postharvest Decay and Diseases

Postharvest diseases caused by microorganisms are a major factor in reducing the quality and shelf life of horticultural crops. Melatonin enhances resistance to pathogens by stimulating plant defense pathways, including the phenylpropanoid (PP) pathway and disease-related proteins (PRs) (Fan et al., 2022). By activating defense enzymes such as PAL, chitinase, and 4-coumaric acid-CoA ligase, this compound can significantly inhibit the occurrence of diseases such as anthracnose, gray mold (*Botrytis cinerea*), and black spot caused by *Alternaria alternata* in crops such as mango, blueberry, and pear (MiaoLian et al., 2022; Wei et al., 2025; Njie et al., 2022). In addition to directly inhibiting pathogens, melatonin prevents physiological damage including browning, frost damage, and tissue deterioration in crops such as tomato and cassava by regulating redox homeostasis and enhancing the antioxidant system (Aghdam et al., 2019; Ma et al., 2016). Overall, melatonin has great potential in managing decay and maintaining the quality of postharvest products with a multifaceted approach.

8. Role of Melatonin in Inhibiting Oxidative Stress

Excessive production of Reactive Oxygen Species (ROS) and subsequent lipid peroxidation are

considered to be the main factors in cell membrane damage and accelerating the aging process in postharvest products (Li et al., 2016). Melatonin can effectively inhibit this destructive process by playing a dual role as a direct antioxidant and a signaling molecule (Shi et al., 2015). Melatonin treatment leads to efficient scavenging of ROS and a significant reduction in the levels of oxidative products, including malondialdehyde (MDA), by inducing the activities of key antioxidant enzymes such as SOD, CAT, APX, POD, and GR, as well as by improving the glutathione to glutathione disulfide ratio (Li et al., 2024; Wan et al., 2024). In addition to protecting membrane integrity and reducing chilling-induced damage, melatonin increases the concentrations of phenolic compounds, anthocyanins, vitamins, and other nutrients by stimulating biosynthetic pathways (such as the shikimate pathway) (Bhardwaj et al., 2022). Extensive evidence supports that melatonin maintains the nutritional quality and significantly increases the shelf life of horticultural products by regulating cellular defense responses and enhancing antioxidant capacity (Borthakur et al., 2024; Kang et al., 2025).

9. Conclusion

Melatonin has emerged as a multifunctional, environmentally benign signaling molecule with considerable potential in the postharvest physiological management of horticultural crops. Converging experimental and molecular evidence indicates that melatonin acts as a central regulatory hub that preserves tissue and membrane integrity, attenuates oxidative damage, and effectively delays ripening and senescence through the modulation of hormonal signaling networks, the reinforcement of plant defense systems, and the stimulation of both enzymatic and non-enzymatic antioxidant machinery. Furthermore, its capacity to enhance tolerance to biotic and abiotic stresses—including chilling and frost injury—and to elevate the concentration of health-promoting bioactive compounds underscores its strategic relevance for safeguarding both the nutritional and commercial value of harvested produce.

However, despite these promising findings, the translation of melatonin-based treatments from controlled laboratory conditions to routine commercial application remains constrained by several unresolved challenges. First, the dose–response relationship of melatonin is frequently non-linear and species-, cultivar-, and maturity-dependent; both suboptimal and supraoptimal concentrations can produce negligible or even detrimental effects, necessitating the precise determination of optimal doses for each commodity and storage scenario. Second, the efficacy of melatonin is strongly influenced by the application strategy—including timing (preharvest vs. postharvest), mode of delivery (dip, spray, fumigation, or incorporation into coatings), carrier formulation, and interaction with storage environment—yet standardized protocols for these variables are still lacking. Third, the economic feasibility of large-scale application, particularly regarding the cost of formulation, labor, and infrastructure, has not been rigorously assessed within real supply-chain contexts. Finally, the absence of validated quality benchmarks, regulatory frameworks, and consumer acceptance data limits the industrial adoption of melatonin-based postharvest interventions.

To address these gaps, future research should be directed along three complementary axes: (i)

elucidation of the precise molecular mechanisms underlying melatonin action through integrated omics approaches and the identification of key regulatory transcription factors; (ii) systematic evaluation of the synergistic effects of melatonin with emerging postharvest technologies, such as edible coatings, modified/controlled atmosphere storage, and biocontrol agents; and (iii) optimization of application protocols—including dose, formulation (e.g., nanoencapsulation), and timing—together with comprehensive techno-economic and life-cycle assessments under pilot and commercial-scale conditions. Ultimately, once these translational barriers are overcome, melatonin can serve as a sustainable, cost-effective, and innovative solution for reducing postharvest losses and enhancing the resilience of the horticultural supply chain.

References

- [1] Abou-Elwafa, S.M. and Safaa, Z., 2025. Effect of melatonin and ascorbic acid application on delay of senescence and keeping quality of fresh broccoli florets. *Horticulture Research Journal*, 3 (2), 1–14.
- [2] Adhikary, T., Gill, P.S., Jawandha, S.K., Bhardwaj, R.D. and Anurag, R.K. 2021. Browning and quality management of pear fruit by salicylic acid treatment during low temperature storage. *Journal of the Science of Food and Agriculture*, 101, 853–862.
- [3] Aghdam, M.S., Jannatizadeh, A., Nojadeh, M.S. and Ebrahimzadeh, A. 2019. Exogenous melatonin ameliorates chilling injury in cut anthurium flowers during low temperature storage. *Postharvest Biology and Technology*, 148, 184–191.
- [4] Aghdam, M.S., Luo, Z., Jannatizadeh, A., Sheikh-Assadi, M., Sharafi, Y., Farmani, B., Fard, J.R. and Razavi, F. 2019. Employing exogenous melatonin applying confers chilling tolerance in tomato fruits by up regulating ZAT2/6/12 giving rise to promoting endogenous polyamines, proline, and nitric oxide accumulation by triggering arginine pathway activity. *Food Chemistry*. 275, 549–556.
- [5] Aghdam, M.S., Luo, Z., Li, L., Jannatizadeh, A., Fard, J.R. and Pirzad, F. 2020. Melatonin treatment maintains nutraceutical properties of

- pomegranate fruits during cold storage. Food Chemistry. 303, 125385.
- [6] Aghdam, M.S. and Fard, J.R. 2017. Melatonin treatment attenuates postharvest decay and maintains nutritional quality of strawberry fruits (*Fragaria x ananassa* cv. Selva) by enhancing GABA shunt activity. Food Chemistry, 221, 1650–1657.
- [7] Aghdam, M.S., Mukherjee, S., Flores, F.B., Arnao, M.B., Luo, Z. and Corpas, F.J. 2022. Functions of Melatonin during Postharvest of Horticultural Crops. Plant and Cell Physiology, 63, 1764–1786.
- [8] Ahmad, I., Song, X., Hussein Ibrahim, M.E., Jamal, Y., Younas, M.U., Zhu, G., Zhou, G. and Adam Ali, A.Y. 2023. The Role of Melatonin in Plant Growth and Metabolism, and Its Interplay with Nitric Oxide and Auxin in Plants under Different Types of Abiotic Stress. Frontiers in Plant Science, 14, 1108507.
- [9] Ahmad, S., Kamran, M., Ding, R., Meng, X., Wang, H., Ahmad, I., Fahad, S. and Han, Q. 2019. Exogenous melatonin confers drought stress by promoting plant growth, photosynthetic capacity and antioxidant defense system of maize seedlings. Peer journal, 7, e7793.
- [10] Arah, I.K., Amaglo, H., Kumah, E.K. and Ofori, H. 2015. Preharvest and postharvest factors affecting the quality and shelf life of harvested tomatoes: a mini review. International Journal of Agronomy, <https://doi.org/10.1155/2015/478041>.
- [11] Arnao, M.B. and Hernández-Ruiz, J. 2018. Melatonin and its relationship to plant hormones. Annals of Botany, 121, 195–207.
- [12] Arnao, M.B. and Hernández-Ruiz, J. 2020. Melatonin in flowering, fruit set and fruit ripening. Plant Reproduction, 33, 77–87.
- [13] Arnao, M.B. and Hernández-Ruiz, J. 2014. Melatonin: Plant Growth Regulator and/or Biostimulator during Stress? Trends in Plant Science, 19, 789–797.
- [14] Badiche-El Hilali, F., Garcia-Pastor, M.E., Valverde, J.M., Castillo, S., Valero, D. and Serrano, M. 2024. Melatonin as an efficient and eco-friendly tool to increase yield and to maintain quality attributes during lemon storage. International Journal of Molecular Sciences, 25 (18).
- [15] Bal, E. 2023. Impact of chitosan-melatonin composite coating on postharvest quality of sweet cherry. Applied Fruit Science, 66 (2), 763–770.
- [16] Baswal, A.K., Dhaliwal, H.S., Singh, Z., Mahajan, B.V.C. and Gill, K.S. 2020. Postharvest application of methyl jasmonate, 1-methylcyclopropene and salicylic acid extends the cold storage life and maintain the quality of ‘Kinnow’ mandarin (*Citrus nobilis* L. X *C. deliciosa* L.) fruit. Postharvest Biology and Technology, 161, 111064.
- [17] Bhardwaj, R., Pareek, S., Mani, S., Domínguez-Avila, J.A. and González-Aguilar, G.A. 2022. A melatonin treatment delays postharvest senescence, maintains quality, reduces chilling injury, and regulates antioxidant metabolism in mango fruit. Journal of Food Quality, 2022, 2379556.
- [18] Borthakur, P., Chinnasamy, K., Paramasivam, S.K., Venkatachalam, S., Alagarswamy, S., Iruthayasamy, J., Thiagarajan, E. and Muthusamy, S. 2024. Exogenous melatonin as pre- and postharvest application on quality attributes, antioxidant capacity, and extension of shelf life of papaya. Horticulturae, 10 (10), 1099.
- [19] Bose, S.K., Howlader, P., Jia, X., Wang, W. and Yin, H. 2019. Alginate oligosaccharide postharvest treatment preserves fruit quality and increase storage life via Absciscic acid signaling in strawberry. Food Chemistry, 283, 665–674.
- [20] Bose, S.K., Howlader, P., Wang, W. and Yin, H. 2021. Oligosaccharide is a promising natural preservative for improving postharvest preservation of fruit: A review. Food Chemistry, 341, 128178.
- [21] Cai, S., Zhang, Z., Wang, J., Fu, Y., Zhang, Z., Khan, M. and Cong, X. 2024. Effect of exogenous melatonin on postharvest storage quality of passion fruit through antioxidant metabolism. LWT, 194, 115835.
- [22] Cao, S., Shao, J., Shi, L., Xu, L., Shen, Z., Chen, W. and Yang, Z. 2018. Melatonin increases chilling tolerance in postharvest peach fruit by alleviating oxidative damage. Scientific Reports, 8(1): 806.
- [23] Cao, J.J. Yu, Z.C., Zhang, Y., Li, B.H., Liang, W.X. and Wang, C.X. 2017. Control efficiency of exogenous melatonin against post-harvest apple grey mold and its influence on the activity of

- defensive enzymes. *Plant Physiology Journal*, 53, 1760.
- [24] Cao, S., Qu, G., Ma, C., Ba, L., Ji, N., Meng, L., Lei, J. and Wang, R. 2022. Effects of melatonin treatment on the physiological quality and cell wall metabolites in kiwifruit. *Food Science & Technology*. (N. Y.) 42.
- [25] Cao, S., Song, C., Shao, J., Bian, K., Chen, W. and Yang, Z. 2016. Exogenous melatonin treatment increases chilling tolerance and induces defense response in harvested peach fruit during cold storage. *Journal of Agricultural and Food Chemistry*, 64, 5215–5222.
- [26] Carrión-Antolí, A., Lorente-Mento, J.M., Valverde, J.M., Castillo, S., Valero, D. and Serrano, M. 2021. Effects of melatonin treatment on sweet cherry tree yield and fruit quality. *Agronomy*, 12(1):3.
- [27] Charoenphun, N., Pham, N.H., Rattanawut, J. and Venkatachalam, K. 2024. Exogenous application of melatonin on the preservation of physicochemical and enzymatic qualities of pepper fruit from chilling injury. *Horticulturae*, 10 (6), 550.
- [28] Chen, H., Lin, H., Jiang, X., Lin, M. and Fan, Z. 2022. Amelioration of chilling injury and enhancement of quality maintenance in cold-stored guava fruit by melatonin treatment. *Food Chemistry X*. 14: 100297.
- [29] Chen, Z., Lin, W., Yang, S., Cui, W., Zhang, S., Peng, Z., Li, Y., Zheng, Y., Xie, F. and Li, M. 2026. Exogenous Melatonin Enhances the Salt Tolerance of Celery (*Apium graveolens* L.) by Regulating Osmotic Adaptation and Energy Metabolism via Starch and Sucrose Metabolic Pathways. *International Journal of Molecular Sciences*, 27, 1299.
- [30] Dai, L., Li, J., Harmens, H., Zheng, X. and Zhang, C. 2020. Melatonin enhances drought resistance by regulating leaf stomatal behaviour, root growth and catalase activity in two contrasting rapeseed (*Brassica napus* L.) genotypes. *Plant Physiology and Biochemistry*, 149: 86–95.
- [31] Devi, M., Jeyakumar, P., Ravichandran, V., Arul, L., Balakrishnan, M. and Kavino, M. 2025. Melatonin application extends banana shelf life by delayed ripening. *Plant Science Today*.
- [32] Eisa, E.A., Tilly-Mándy, A. and Honfi, P. 2026. Phytomelatonin in Ornamental Horticulture: A Comprehensive Review of Growth Promotion, Stress Tolerance, and Post-Harvest Quality. *International Journal of Molecular Sciences*, 27 (4): 1645.
- [33] El-Naby, A., Abdelkhalek Ahmed Mohamed, M. and El-Naggar, Y.I.M. 2019. Effect of melatonin, GA3 and NAA on vegetative growth, yield and quality of ‘Canino’apricot fruits. *Acta Scientiarum Polonorum. Hortorum Cultus*, 18 (3): 167–74.
- [34] Ema, N.R., Mithu, M.A.H. and Sayem, A. 2024. Exploring driving factors in employing waste reduction tools to alleviate the global food security and sustainability. *Heliyon*, 10, e28192.
- [35] Fan, A., Wan, C., Liu, H., Xiong, X., Nong, Y., Kahramanoglu, I., Yang, R. and Zeng, L. 2023. Melatonin treatment maintains the quality and delays senescence of postharvest cattails (*Typha latifolia* L.) during cold storage. *Food Chemistry: X*, 19, 100796.
- [36] Fan, S., Li, Q., Feng, S., Lei, Q., Abbas, F., Yao, Y., Chen, W., Li, X. and Zhu, X. 2022. Melatonin maintains fruit quality and reduces anthracnose in postharvest papaya via enhancement of antioxidants and inhibition of pathogen development. *Antioxidants*, 11, 804.
- [37] Fan, S., Xiong, T., Lei, Q., Tan, Q., Cai, J., Song, Z., Yang, M., Chen, W., Li, X. and Zhu, X. 2022. Melatonin treatment improves postharvest preservation and resistance of guava fruit (*Psidium guajava* L.). *Foods*, 11, 262.
- [38] Favero, G., Sulas, F., Labanca, M., Scilla, F., Punzi, C.F., Lonati, C. and Rezzani, R. 2026. Glioblastoma and Melatonin's Effects, A Narrative Review. *Cancers*, 18, 703.
- [39] Ferreira, C., Ribeiro, C. and Nunes, F.M. 2024. Effect of storage conditions on phenolic composition, vitamin C and antioxidant activity of ‘Golden Delicious’ and ‘Red Delicious’ apples. *Postharvest Biology and Technology*, 210, 112754.
- [40] Fracassetti, D., Vigentini, I., Lo Faro, A. F. F., De Nisi, P., Foschino, R., Tirelli, A., Orioli, M., and Iriti, M. 2019. Assessment of Tryptophan, Tryptophan Ethylester, and Melatonin Derivatives in Red Wine by SPE-HPLC-FL and SPE-HPLC-MS Methods. *Foods*, 8(3), 99.
- [41] Galano, A. and Reiter, R.J. 2018. Melatonin and its metabolites vs oxidative stress, From

- individual actions to collective protection. Journal of Pineal Research, 65, e12514.
- [42] Gao, H., Lu, Z., Yang, Y., Wang, D., Yang, T., Cao, M. and Cao, W. 2018. Melatonin treatment reduces chilling injury in peach fruit through its regulation of membrane fatty acid contents and phenolic metabolism. Food Chemistry, 245, 659–666.
- [43] Gao, H., Zhang, Z.K., Chai, H.K., Cheng, N., Yang, Y., Wang, D.N., Yang, T. and Cao, W. 2016. Melatonin Treatment Delays Postharvest Senescence and Regulates Reactive Oxygen Species Metabolism in Peach Fruit. Postharvest Biology and Technology, 118, 103–110.
- [44] Gao, T., Liu, X., Tan, K., Zhang, D., Zhu, B., Ma, F. and Li, C. 2022. Introducing Melatonin to the Horticultural Industry: Physiological Roles, Potential Applications, and Challenges. Horticulture Research, 9, uhac094.
- [45] Ghorbani, B. and Najafzadeh, R. 2022. The Role of Melatonin on postharvest properties of sweet cherry fruit cv. siah mashhad during storage. Journal of Horticultural Science, 36 (1), 71–85.
- [46] Gökbayrak, Z., Engin, H. and Kiraz, H. 2020. Effects of melatonin and IAA on adventitious root formation in rootstock 5BB and cv. cabernet sauvignon (*Vitis vinifera* L.). Kahramanmaraş Sütçü İmam Üniversitesi Tarım Ve Doğa Dergisi, 23 (4): 835–41.
- [47] Gurjar, P.S., Berwal, M.K., Kumar, R., Sarolia, D.K. and Choudhary, M.K. 2024. Post-harvest melatonin treatment maintains freshness and delayed peel browning by sustaining higher antioxidants in Indian jujube (*ziziphus mauritiana* lamk) fruits. Applied Fruit Science. 67 (1).
- [48] Gutiérrez-Villamil, D.A., Balaguera-López, H.E. and Álvarez-Herrera, J.G. 2023. Brassinosteroids Improve Postharvest Quality, Antioxidant Compounds, and Reduce Chilling Injury in ‘Arrayana’ Mandarin Fruits under Cold Storage. Horticulturae, 9, 622.
- [49] Hanci, F. and Hasan, A.H.H. 2025. Acute and Delayed Effects of Melatonin Pretreatment Against Cold Stress in Leek (*Allium ampeloprasum* L. var. porrum). Horticulturae, 11, 1208.
- [50] Hanif, A., Ahmad, S., Shahzad, S., Liaquat, M. and Anwar, R. 2020. Postharvest application of salicylic acid reduced decay and enhanced storage life of papaya fruit during cold storage. Journal of Food Measurement and Characterization, 14, 3078–3088.
- [51] He, Z., Wen, C. and Xu, W. 2023. Effects of Endogenous Melatonin Deficiency on the Growth, Productivity, and Fruit Quality Properties of Tomato Plants. Horticulturae, 9, 851.
- [52] Hei, H., Tang, H., Zhao, R., Li, G. and Shi, F. 2025. Effect of Melatonin Treatment on storage quality and antioxidant system of postharvest winter jujube (*Zizyphus jujube* mill. cv. dongzao). Foods, 14 (4), 576.
- [53] Hernández-Ruiz, J., Ruiz-Cano, D., Giraldo-Acosta, M., Cano, A. and Arnao, M.B. 2022. Melatonin in Brassicaceae, Role in Postharvest and Interesting Phytochemicals. Molecules, 27, 1523.
- [54] Hu, W., Kong, H., Guo, Y., Zhang, Y., Ding, Z., Tie, W., Yan, Y., Huang, Q., Peng, M., Shi, H. and Guo, A. 2016. Comparative Physiological and Transcriptomic Analyses Reveal the Actions of Melatonin in the Delay of Postharvest Physiological Deterioration of Cassava. Frontiers in Plant Science. 7: 736.
- [55] Hu, C., Zheng, Y., Tong, C. and Zhang, D. 2022. Effects of exogenous melatonin on plant growth, root hormones and photosynthetic characteristics of Trifoliate orange subjected to salt stress. Plant Growth Regulation, 97, 551–558.
- [56] Hu, S., Ma, Y., Xie, B., Hou, Y., Jia, Z., Zhao, L., Zheng, Y. and Jin, P. 2022. 24-Epibrassinolide improves chilling tolerance by regulating PpCBF5-mediated membrane lipid metabolism in peach fruit. Postharvest Biology and Technology, 186, 111844.
- [57] Hu, W., Tie, W., Ou, W., Yan, Y., Kong, H., Zuo, J., Ding, X., Ding, Z., Liu, Y., Wu, C., Guo, Y., Shi, H., Li, K. and Guo, A. 2018. Crosstalk between calcium and melatonin affects postharvest physiological deterioration and quality loss in cassava. Postharvest Biology and Technology, 140, 42–49.
- [58] Hu, W., Yang, H., Tie, W., Yan, Y., Ding, Z., Liu, Y., Wu, C., Wang, J., Reiter, R.J., Tan, D.X., Shi, H., Xu, B. and Jin, Z. 2017. Natural Variation in Banana Varieties Highlights the Role of Melatonin in Postharvest Ripening and Quality. Journal of Agricultural and Food Chemistry, 65(46): 9987-9994.

- [59] Huang, N.C. and Wong, C.S. 2026. Melatonin as an Integrative Adjunct in Multimodal Analgesia, Linking Circadian Regulation, Anti-Inflammatory Modulation, and Opioid-Sparing Mechanisms. *International Journal of Molecular Sciences*, 27, 2046.
- [60] Huang, T., Kou, X., Qiao, L., Li, J., Luo, D., Wang, X. and Cao, S. 2024. Maintaining quality of postharvest green pepper fruit using melatonin by regulating membrane lipid metabolism and enhancing antioxidant capacity. *Food Chemistry*, 460, 140671.
- [61] Islam, M., Ali, S., Nawaz, A., Naz, S., Ejaz, S., Shah, A.A. and Razzaq, K. 2022. Postharvest 24-epibrassinolide treatment alleviates pomegranate fruit chilling injury by regulating proline metabolism and antioxidant activities. *Postharvest Biology and Technology*, 188, 111906.
- [62] Jabbari, M. and Jabbari, M. 2023. Regulatory and protective role of plant melatonin. *Iranian Journal of Biology*, 7 (autumn and winter), 179-193.
- [63] Jannatizadeh, A., Aghdam, M.S., Luo, Z. and Razavi, F. 2019. Impact of exogenous melatonin application on chilling injury in tomato fruits during cold storage. *Food and Bioprocess Technology*, 12 (5), 741–750.
- [64] Jia Zhiwei, S.M., Jinmei, C., Kanghua, S. and Lubin, Z. 2019. Effect of melatonin treatment on the preservation of the flower stalk of brassica campestris L. ssp. chinesis var. utilisen et lee after. *Chinese Journal of Tropical Crops*, 40 (7), 1413–1420.
- [65] Juhnevica-Radenkova, K., Moreno, D.A., Ikase, L., Drudze, I. and Radenkovs, V. 2020. Naturally occurring melatonin: Sources and possible ways of its biosynthesis. *Comprehensive Reviews in Food Science and Food Safety*. 19(6): 4008–30.
- [66] Kahramanog˘lu, I'. 2024. Melatonin prolongs the postharvest storability of strawberry fruits through reduced oxidative stress and improved enzyme activity. *Hortscience*, 59 (9), 1350–1357.
- [67] Kang, D., Zeng, J., Lu, D., Yue, L., Zhong, M., Kang, Y., Guo, J. and Yang, X. 2025. Melatonin maintains postharvest quality by modulating the metabolisms of cell wall and sugar in flowering Chinese cabbage. *Scientific Reports*, 15 (1), 5396.
- [68] Küçüker, E., Kuru Berk, S., Melda Çolak, A. and Gundogdu, M. 2024. Melatonin use in post-harvest fruit physiology: effect of melatonin treatments on quality properties, primer and seconder metabolites contents of hawthorn fruit during cold storage. *Journal of Plant Growth Regulation*, 44 (5), 2023–2036.
- [69] Kumar Bose, S. and Howlade, P. 2026. Melatonin: An eco-friendly biomolecule for maintaining postharvest quality and enhancing storage life of horticultural crops. *Journal of Experimental Horticulture*, 1:100003.
- [70] Kumari, A., Singh, S.K., Mathpal, B., Verma, K.K., Garg, V.K., Bhattacharyya, M. and Bhatt, R. 2023. The Biosynthesis, Mechanism of Action, and Physiological Functions of Melatonin in Horticultural Plants, A Review. *Horticulturae*, 9, 913.
- [71] Kumari, A., Singh, S.K. and Rai, G.K. 2021. Physiological and Biochemical Response of Plants under Drought Stress. In *Plant Abiotic Stress Tolerance: Physiochemical and Molecular Avenues*; Deepika Book Agency: Delhi, India, pp. 77–92.
- [72] Lau, K.Q., Sabran, M.R. and Shafie, S.R. 2021. Utilization of Vegetable and Fruit By-products as Functional Ingredient and Food. *Frontiers in Nutrition*. 8, 661693.
- [73] Lei, Q., Wang, L., Tan, D.X., Zhao, Y., Zheng, X.D., Chen, H., Li, Q.T., Zuo, B.X. and Kong, J. 2013. Identification of genes for melatonin synthetic enzymes in 'Red Fuji' apple (*Malus domestica* Borkh. cv. Red) and their expression and melatonin production during fruit development. *Journal of Pineal Research*, 55(4): 443-51.
- [74] Li, H., He, J., Yang, X., Li, X., Luo, D., Wei, C., Ma, J., Zhang, Y., Yang, J. and Zhang, X. 2016. Glutathione-dependent induction of local and systemic defense against oxidative stress by exogenous melatonin in cucumber (*Cucumis sativus* L.). *Journal of Pineal Research*, 60(2): 206-16.
- [75] Li, J., Liu, J., Zhu, T., Zhao, C., Li, L. and Chen, M. 2019. The Role of Melatonin in Salt Stress Responses. *International Journal of Molecular Sciences*, 20, 1735.
- [76] Li, J.X., Luo, M.M., Tong, C.L., Zhang, D.J. and Zha, Q. 2023. Advances in fruit coloring research

- in grapevine, an overview. *Plant Growth Regulation*, 103, 51–63.
- [77] Li, P., Zhang, R., Zhou, H., Mo, Y., Wu, S., Zhang, X., Xie, Z., Zhang, T., Zhao, K., Lv, J. and Deng, M. 2024. Melatonin delays softening of postharvest pepper fruits (*Capsicum annuum* L.) by regulating cell wall degradation, membrane stability and antioxidant systems. *Postharvest Biology and Technology*, 212, 112852.
- [78] Li, S., Xu, Y., Bi, Y., Zhang, B., Shen, S., Jiang, T. and Zheng, X. 2019. Melatonin treatment inhibits gray mold and induces disease resistance in cherry tomato fruit during postharvest. *Postharvest Biology and Technology*, 157, 110962.
- [79] Li, X., Xu, D., Xu, H. and Cao, P. 2026. Melatonin Supplementation in Sex-Sorted Nili-Ravi Buffalo Semen, Effect on Sperm Quality, Subsequent in Vitro Embryo Development, and Pregnancy Outcomes. *Antioxidants*, 15, 344.
- [80] Lin, X., Wang, L., Hou, Y., Zheng, Y. and Jin, P. 2020. A combination of melatonin and ethanol treatment improves postharvest quality in bitter melon fruit. *Foods*, 9, 1376.
- [81] Lin, Y., Fan, L., Xia, X., Wang, Z., Yin, Y., Cheng, Y. and Li, Z. 2019. Melatonin decreases resistance to postharvest green mold on citrus fruit by scavenging defense-related reactive oxygen species. *Postharvest Biology and Technology*, 153, 21–30.
- [82] Liu, S., Huang, H., Huber, D.J., Pan, Y., Shi, X. and Zhang, Z. 2020. Delay of ripening and softening in ‘Guifei’ mango fruit by postharvest application of melatonin. *Postharvest Biology and Technology*, 163: 111136.
- [83] Liu, C., Zheng, H., Sheng, K., Liu, W. and Zheng, L. 2018. Effects of melatonin treatment on the postharvest quality of strawberry fruit. *Postharvest Biology and Technology*, 139, 47–55.
- [84] Liu, D.D., Sun, X.S., Liu, L., Shi, H.D., Chen, S.Y. and Zhao, D.K. 2019. Overexpression of the melatonin synthesis-related gene SICOMT1 improves the resistance of tomato to salt stress. *Molecules*, 24 (8), 1514.
- [85] Liu, J., Yang, J., Zhang, H., Cong, L., Zhai, R., Yang, C., Wang, Z., Ma, F. and Xu, L. 2019. Melatonin inhibits ethylene synthesis via nitric oxide regulation to delay postharvest senescence in pears. *Journal of Agricultural and Food Chemistry*, 67 (8), 2279–2288.
- [86] Ma, Q., Zhang, T., Zhang, P. and Wang, Z.Y. 2016. Melatonin attenuates postharvest physiological deterioration of cassava storage roots. *Journal of Pineal Research*, 60 (4), 424–434.
- [87] Mandal, M.K., Suren, H., Ward, B., Boroujerdi, A. and Kousik, C. 2018. Differential roles of melatonin in plant-host resistance and pathogen suppression in cucurbits. *Journal of Pineal Research*, 65 (3), e12505.
- [88] Mao, J., Niu, C., Li, K., Chen, S., Tahir, M.M., Han, M. and Zhang, D. 2020. Melatonin promotes adventitious root formation in apple by promoting the function of MdWOX11. *BMC Plant Biology*, 20(1):1–11.
- [89] Massolo, J.F., Sánchez, R., Zaro, M.J., Concellón, A. and Vicente, A.R. 2022. Low-dose prestorage 24-epibrassinolide spray enhances postharvest chilling tolerance in zucchini squash (*Cucurbita pepo* L.) by eliciting peroxidase and phenolic antioxidants. *Journal of Food Processing and Preservation*, 46, e16576.
- [90] Medina-Santamarina, J., Serrano, M., Lorente-Mento, J.M., García-Pastor, M.E., Zapata, P.J., Valero, D. and Guillén, F. 2021. Melatonin treatment of pomegranate trees increases crop yield and quality parameters at harvest and during storage. *Agronomy*, 11(5): 861.
- [91] Meng, J.F., Yu, Y., Shi, T.C., Fu, Y.S., Zhao, T. and Zhang, Z.W. 2019. Melatonin treatment of perversion grape berries modifies phenolic components and antioxidant activity of grapes and wine. *Journal of Food Science and Technology*, 39 (1), 35–42.
- [92] MiaoLian, X., Fan, W., Cheng, L.S., YinBao, W., LiuHua, X., WenWen, P., JinYin, C. and Ming, C. 2022. Effects of melatonin treatment on resistance to black spot and postharvest storage quality of pear fruit. *Scient. Agril. Sinic.* 55 (4), 785–795.
- [93] Miranda, S., Vilches, P., Suazo, M., Pavez, L., García, K., Méndez, M.A., González, M., Meisel, L.A., Defilippi, B.G. and del Pozo, T. 2020. Melatonin triggers metabolic and gene expression changes leading to improved quality traits of two sweet cherry cultivars during cold storage. *Food Chemistry*, 319, 126360.

- [94] Nguyen, T. and Ha, S. 2025. Melatonin: An Eco-Friendly Preservative for Improving Post-Harvest Quality and Longevity of Cut Flowers. *Horticulturae*, 11(6), 574.
- [95] Nitu, N.J., Ullah, M.S., Howlader, P., Mehedi, M.N.H., Meem, H.Z. and Bose, S.K. 2024. Chitosan oligosaccharides maintained postharvest quality and increased shelf life of mango. *Journal of Horticulture and Postharvest Research*. 8 (1), 43–66.
- [96] Njie, A., Zhang, W., Dong, X., Lu, C., Pan, X. and Liu, Q. 2022. Effect of melatonin on fruit quality via decay inhibition and enhancement of ant oxidative enzyme activities and genes expression of two mango cultivars during cold storage. *Foods*, 11 (20).
- [97] Onik, J.C., Wai, S.C., Li, A., Lin, Q., Sun, Q., Wang, Z. and Duan, Y. 2021. Melatonin treatment reduces ethylene production and maintains fruit quality in apple during postharvest storage. *Food Chemistry*, 337, 127753.
- [98] Orsini, F., Kahane, R., Nono-Womdim, R. and Gianquinto, G. 2013. Urban Agriculture in the Developing World: A Review. *Agronomy for Sustainable Development*, 33, 695–720.
- [99] Padilla-González, P.A., Garrido-Auñón, F., García-Pastor, M.E., Guillén, F., Serrano, M., Valero, D. and Agulló, V. 2026. Melatonin as a Pre-and Postharvest Tool for Enhancing Fruit Quality. *Plants*, 15, 331.
- [100] Paroni, R., Dei Cas, M., Rizzo, J., Ghidoni, R., Montagna, M.T., Rubino, F.M. and Iriti, M. 2019. Bioactive phytochemicals of tree nuts. Determination of the melatonin and sphingolipid content in almonds and pistachios. *Journal of Food Composition and Analysis*, 82: 103227.
- [101] Patil, M., Sharma, S., Sridhar, K., Anurag, R.K., Grover, K., Dharni, K., Mahajan, S. and Sharma, M. 2024. Effect of postharvest treatments and storage temperature on the physiological, nutritional, and shelf-life of broccoli (*Brassica oleracea*) microgreens. *Scientia Horticulturae*, 327, 112805.
- [102] Ping, X., Huang, T., Shiqi, L., Meijiao, H., Zhaoyin, G., Jialiang, L. and Zhang, Z. 2024. Effect of melatonin on chilling injury and ripening of postharvest mango fruits during shelf life after refrigeration and underlying physiological mechanism. *Food scientists*, (N. Y.) 45 (8), 218–227.
- [103] Poveda, J. 2020. Use of plant-defense hormones against pathogen-diseases of postharvest fresh produce. *Physiological and Molecular Plant Pathology*, 111, 101521.
- [104] Pranil, T., Moongngarm, A., Manwiwattanakul, G., Loypimai, P. and Kerr, W.L. 2021. Melatonin and its derivative contents in tropical fruits and fruit tablets. *Journal of Food Composition and Analysis*, 103: 104109.
- [105] Qi, Z.Y., Wang, K.X., Yan, M.Y., Kanwar, M., Li, D.Y., Wijaya, L., Alyemeni, M., Ahmad, P. and Zhou, J. 2018. Melatonin Alleviates High Temperature-Induced Pollen Abortion in *Solanum lycopersicum*. *Molecules*, 23, 386.
- [106] Qu, G., Ba, L., Wang, R., Li, J., Ma, C., Ji, N. and Cao, S. 2022. Effects of melatonin on blueberry fruit quality and cell wall metabolism during low temperature storage. *Food Science & Technology*, 42.
- [107] Raak, N., Symmank, C., Zahn, S., Aschemann-Witzel, J. and Rohm, H. 2017. Processing and product related causes for food waste and implications for the food supply chain. *Waste Management*, 61, 461–472.
- [108] Rahmanzadeh-Ishkeh, S., Shirzad, H., Tofighi, Z., Fattahi, M. and Ghosta, Y. 2024. Exogenous melatonin prolongs raspberry postharvest life quality by increasing some antioxidant and enzyme activity and phytochemical contents. *Scientific Reports*, 14 (1), 11508.
- [109] Rastegar, S., Hassanzadeh Khankahdani, H. and Rahimzadeh, M. 2020. Effects of melatonin treatment on the biochemical changes and antioxidant enzyme activity of mango fruit during storage. *Scientia Horticulturae*, 259, 108835.
- [110] Riva, S.C., Opara, U.O. and Fawole, O.A. 2020. Recent developments on postharvest application of edible coatings on stone fruit: A review. *Scientia Horticulturae*, 262, 109074.
- [111] Salehi, B., Sharopov, F., Fokou, P.V.T., Kobylinska, A., Jonge, L., Tadio, K., Sharifi-Rad, J., Posmyk, M.M., Martorell, M., Martins, N. and Iriti, M. 2019. Melatonin in Medicinal and Food Plants: Occurrence, Bioavailability, and Health Potential for Humans. *Cells*, 8(7): 681.

- [112] Saud, S., Jiang, Z., Chen, S. and Fahad, S. 2023. Interaction of melatonin on post-harvest physiology and quality of horticultural crops. *Scientia Horticulturae*, 321, 112286.
- [113] Seyed, M.H.M., Somayeh, R., Valiollah, G.O. and Orang, K. 2020. Ameliorative effect of melatonin against storage chilling injury in pomegranate husk and arils through promoting the antioxidant system. *Scientia Horticulturae*, 295, 110889.
- [114] Shang, F., Liu, R., Wu, W., Han, Y., Fang, X., Chen, H. and Gao, H. 2021. Effects of melatonin on the components, quality and antioxidant activities of blueberry fruits. *LWT*, 147, 111582.
- [115] Shao, R., Wang, Y., He, C. and Chen, L. 2024. Melatonin and its emerging physiological role in reproduction: a review and update. *Current Molecular Medicine*, 24: 449–56.
- [116] Sharafi, Y., Aghdam, M.S., Luo, Z., Jannatizadeh, A., Razavi, F., Fard, J.R. and Farmani, B. 2019. Melatonin treatment promotes endogenous melatonin accumulation and triggers GABA shunt pathway activity in tomato fruits during cold storage. *Scientia Horticulturae*, 254, 222–227.
- [117] Sharafi, Y., Jannatizadeh, A., Fard, J.R. and Aghdam, M.S. 2021. Melatonin treatment delays senescence and improves antioxidant potential of sweet cherry fruits during cold storage. *Scientia Horticulturae*, 288, 110304.
- [118] Sharma, A. and Zheng, B. 2019. Melatonin mediated regulation of drought stress: Physiological and molecular aspects. *Plants*, 8, 190.
- [119] Shen, T., Liang, D., Wang, X., Lv, X., Wang, J. and Xia, H. 2019. Effect of Melatonin on the Breaking Dormancy of Kiwifruit Seeds. In: IOP conference series: earth and environmental science, vol. 237, No. 5, p. 052073. IOP Publishing.
- [120] Shi, G., Zhou, X., Tong, C. and Zhang, D. 2024. The Physiological and Molecular Mechanisms of Fruit Cracking Alleviation by Exogenous Calcium and GA3 in the Late Navel Orange. *Horticulturae*, 10, 1283.
- [121] Shi, H., Tan, D.X., Reiter, R.J., Ye, T., Yang, F. and Chan, Z. 2015. Melatonin induces class A1 heat-shock factors (HSFA 1s) and their possible involvement of thermo tolerance in Arabidopsis. *Journal of Pineal Research*, 58 (3), 335–342.
- [122] Shi, L., Chen, Y., Dong, W., Li, S., Chen, W., Yang, Z. and Cao, S. 2024. Melatonin delayed senescence by modulating the contents of plant signaling molecules in postharvest okras. *Frontiers in Plant Science*, 15, 1304913.
- [123] Sun, H.J., Luo, M.L., Zhou, X., Zhou, Q. and Ji, S.J. 2020. Influence of melatonin treatment on peel browning of cold-stored 'Nangu' Pears. *Food and Bioprocess Technology*, 13, 1478–1490.
- [124] Tijero, V., Munoz, P. and Munne-Bosch, S. 2019. Melatonin as an inhibitor of sweet cherries ripening in orchard trees. *Plant Physiology and Biochemistry*, 140: 88–95.
- [125] Tongtong, L., Tianhong, P., Ji, M., Hui, L., Xiaoting, Q., Fangfang, H. and Changxia, D. 2018. Melatonin for cut flower preservation with *Rosa hybrida* 'Corolla'. *Zhejiang Agriculture and Forestry University*, 35 (5), 981–986.
- [126] Tu, V.T.C., Tra, H.T.T., An, H.D., Thanh, L.D., Hoang, T.M.N. and Hue, T.T. 2022. Influence of exogenous application of melatonin on biochemical changes of avocado fruit during ripening. *Academia Journal of Biology*, 44 (2), 123–132.
- [127] Uğur, Y. 2023. Extraction and quantification of melatonin in cornelian cherry (*Cornus mas* L.) by ultra-fast liquid chromatography coupled to fluorescence detector (UFLC-FD). *Acta Chromatogr*, 35 (3): 219–26.
- [128] Ullah, S.K., Nitu, N.J., Howlader, P., Mehedi, N.H. and Bose, S.K. 2025. Natural preservatives maintained postharvest quality, reduced decay percentage and increased shelf life of mango. *International Journal of Horticultural Science and Technology*, 1261–1280.
- [129] Umamaheswari, R., Prasannakumar, N.R., Sriram, S., Sharma, S.K., Rao, M.S. and Chaya, M.K. 2020. Biotic Stress Management in Horticultural Crops Using Microbial Intervention. In *Rhizosphere Microbes*; Sharma, S.K., Singh, U.B., Sahu, P.K., Singh, H.V., Sharma, P.K., Eds.; Springer: Singapore, Volume 23, pp. 619–654. ISBN 9789811591532.
- [130] Usanmaz, S., Aşkın, M.A. and Kahramanoğlu, İ. 2022. Effects of foliar melatonin application on the flowering and fruit

- set of pomegranate (*Punica granatum* L. Wonderful). In: V international symposium on pomegranate and minor Mediterranean fruits. 1349. p. 63–72.
- [131] Verde, A., Míguez, J.M. and Gallardo, M. 2019. Melatonin and related bioactive compounds in commercialized date palm fruits (*Phoenix dactylifera* L.): correlation with some antioxidant parameters. *European Food Research and Technology*, 245: 51–9.
- [132] Verde, A., Míguez, J.M. and Gallardo, M. 2022. Role of melatonin in apple fruit during growth and ripening: possible interaction with ethylene. *Plants*, 11(5): 688.
- [133] Verde, A., Míguez, J.M., Leao-Martins, J.M., Gago-Martínez, A. and Gallardo, M. 2022. Melatonin content in walnuts and other commercial nuts Influence of cultivar, ripening and processing (roasting). *Journal of Food Composition and Analysis*, 105: 104180.
- [134] Verde, A., Míguez, J.M. and Gallardo, M. 2022. Role of melatonin in apple fruit during growth and ripening: possible interaction with ethylene. *Plants*, 11 (5): 688.
- [135] Wan, J., Wu, Y., Tong, Z., Su, W., Lin, H. and Fan, Z. 2024. Melatonin treatment alleviates chilling injury of loquat fruit via modulating ROS metabolism. *Foods*, 13 (19): 3050.
- [136] Wang, F., Zhang, X., Yang, Q. and Zhao, Q. 2019. Exogenous melatonin delays postharvest fruit senescence and maintains the quality of sweet cherries. *Food Chemistry*, 301: 125311.
- [137] Wang, M., Li, Y., Li, C., Xu, H., Sun, T. and Ge, Y. 2023. Melatonin induces resistance against *Penicillium expansum* in apple fruit through enhancing phenylpropanoid metabolism. *Physiological and Molecular Plant Pathology*, 127: 102082.
- [138] Wang, D., Randhawa, M.S., Azam, M., Liu, H., Ejaz, S., Ilahy, R., Qadri, R., Khan, M.I., Umer, M.A., Khan, M.A. and Wang, K. 2022. Exogenous melatonin treatment reduces postharvest senescence and maintains the quality of papaya fruit during cold storage. *Frontiers in Plant Science*, 13, 1039373.
- [139] Wang, F., Zhang, X., Yang, Q. and Zhao, Q. 2019. Exogenous melatonin delays postharvest fruit senescence and maintains the quality of sweet cherries. *Food Chemistry*, 301, 125311.
- [140] Wang, H., Zhang, W., Xu, M. and Su, L. 2026. Melatonin-Induced Leaf Growth in *Lithocarpus litseifolius*: A Synergistic Interplay among Hormone Homeostasis, Photosynthetic Enhancement, and Transcriptional Regulation. *Horticulturae*, 12, 694.
- [141] Wang, K., Xing, Q., Ahammed, G.J. and Zhou, J. 2022. Functions and Prospects of Melatonin in Plant Growth, Yield, and Quality. *Journal of Experimental Botany*, 73, 5928–5946.
- [142] Wang, M., Xu, J., Ding, Z. and Xie, J. 2023. Prolong the Postharvest Shelf Life of Spinach through the Ant oxidative Ability of Melatonin. *Food Chemistry*, X, 19, 100769.
- [143] Wang, S., Shi, X., Wang, R., Wang, H., Liu, F. and Laborda, P. 2020. Melatonin in fruit production and postharvest preservation, A review. *Food Chemistry*, 320, 126642.
- [144] Wei, Q., Xiao, L., Du, Q., Zhang, P., Deng, J. and Wang, F. 2025. Postharvest application of melatonin suppresses gray mold in blueberry fruits by regulating cell wall metabolism. *ES Food & Agroforestry*. doi:10.30919/faf1924.
- [145] Xia, H., Shen, Y., Deng, H., Wang, J., Lin, L., Deng, Q., Lv, X., Liang, D., Hu, R., Wang, Z. and Xiong, B. 2021. Melatonin application improves berry coloration, sucrose synthesis, and nutrient absorption in 'Summer Black' grape. *Food Chemistry*, 356: 129713.
- [146] Xie, J., Qin, Z., Pan, J., Li, J., Li, X., Khoo, H.E. and Dong, X. 2022. Melatonin treatment improves postharvest quality and regulates reactive oxygen species metabolism in "Feizixiao" litchi based on principal component analysis. *Frontiers in Plant Science*, 13, 965345.
- [147] Xin, D., Si, J. and Kou, J. 2017. Postharvest exogenous melatonin enhances quality and delays the senescence of cucumber. *Acta Horticulturae*, 44 (5), 891–901.
- [148] Xu, L., Yue, Q., Xiang, G., Bian, F.e. and Yao, Y. 2018. Melatonin promotes ripening of grape berry via increasing the levels of ABA, H₂O₂, and particularly ethylene. *Horticulture Research*, 5, 41.
- [149] Xu, T., Chen, Y. and Kang, H. 2019. Melatonin is a potential target for improving post-harvest preservation of fruits and vegetables. *Frontiers in Plant Science*, 10, 1388.
- [150] Yang, Y., Cui, X. and Zhong, L. 2023. Effect of melatonin on ripening of 'Pearl' guava fruit at

- ambient temperature. Emirates Journal of Food and Agriculture, 35 (1): 32–39.
- [151] Yılmaz, U.Ç.U.R. 2021. Investigation of Melatonin content and antioxidant capacity in grape berries. İnönü Üniversitesi Sağlık Hizmetleri Meslek Yüksek Okulu Dergisi. 9(3): 820–30.
- [152] Zhao, Y., Tan, D.X., Lei, Q., Chen, H., Wang, L., Li, Q.T., Gao, Y. and Kong, J. 2013. Melatonin and its potential biological functions in the fruits of sweet cherry. Journal of Pineal Research, 55 (1): 79–88.
- [153] Ze, Y., Gao, H., Li, T., Yang, B. and Jiang, Y. 2021. Insights into the roles of melatonin in maintaining quality and extending shelf life of postharvest fruits. Trends in Food Science & Technology, 109, 569–578.
- [154] Zhai, R., Liu, J., Liu, F., Zhao, Y., Liu, L., Fang, C., Wang, H., Li, X., Wang, Z., Ma, F. and Xu, L. 2018. Melatonin limited ethylene production, softening and reduced physiology disorder in pear (*Pyrus communis* L.) fruit during senescence. Postharvest Biology and Technology, 139, 38–46.
- [155] Zhang, H., Liu, X., Chen, T., Ji, Y., Shi, K., Wang, L., Zheng, X. and Kong, J. 2018. Melatonin in Apples and Juice: Inhibition of Browning and Microorganism Growth in Apple Juice. Molecules, 23(3), 521.
- [156] Zhang, H., Wang, L., Shi, K., Shan, D., Zhu, Y., Wang, C., Bai, Y., Yan, T., Zheng, X. and Kong, J. 2019. Apple tree flowering is mediated by low level of melatonin under the regulation of seasonal light signal. Journal of Pineal Research, 66 (2): e12551.
- [157] Zhang, L., Yu, Y., Chang, L., Wang, X. and Zhang, S. 2022. Melatonin enhanced the disease resistance by regulating reactive oxygen species metabolism in post-harvest jujube fruit. Journal of Food Processing and Preservation, 46(3): e16363.
- [158] Zhang, T., Wang, J., Sun, Y., Zhang, L. and Zheng, S. 2022. Versatile roles of melatonin in growth and stress tolerance in plants. Journal of Plant Growth Regulation, 41: 1–17.
- [159] Zhang, Y., Gong, Y., Chen, L., Peng, Y., Wang, Q. and Shi J. 2019. Hypotaurine delays senescence of peach fruit by regulating reactive oxygen species metabolism. Scientia Horticulturae, 253: 295–302.
- [160] Zhang, M., Yang, X., Yin, C., Lin, X., Liu, K., Zhang, K., Su, Y., Zou, X., Liao, L., Wang, X., He, S., He, R., Sun, G., He, J., Xiong, B. and Wang, Z. 2024. Effect of exogenous melatonin on antioxidant properties and fruit softening of ‘Fengtang’ plum fruit (*Prunus salicina* Lindl.) during storage at room temperature. Frontiers in Plant Science, 15, 1348744.
- [161] Zhang, Y. 2024. Post-harvest cold shock treatment enhanced antioxidant capacity to reduce chilling injury and improves the shelf life of guava (*Psidium guajava* L.). Frontiers in Sustainable Food Systems, 8, 1297056.
- [162] Zhang, Y., Huber, D.J., Hu, M., Jiang, G., Gao, Z., Xu, X., Jiang, Y. and Zhang, Z. 2018. Delay of postharvest browning in litchi fruit by melatonin via the enhancing of antioxidative processes and oxidation repair. Journal of Agricultural and Food Chemistry, 66, 7475–7484.
- [163] Zheng, H. Liu, W., Liu, S., Liu, C. and Zheng, L. 2019. Effects of melatonin treatment on the enzymatic browning and nutritional quality of fresh-cut pear fruit. Food Chemistry, 299, 125116.
- [164] Zhou, K., Cheng, Q., Dai, J., Liu, Y., Liu, Q., Li, R., Wang, J., Hu, R. and Lin, L. 2023. Effects of exogenous melatonin on sugar and organic acid metabolism in early-ripening peach fruits. PLoS One, 18 (10): e0292959.
- [165] Zhou, K., Li, Y., Hu, L., Zhang, J., Yue, H., Yang, S., Liu, Y., Gong, X. and Ma, F. 2021. Overexpression of MdASMT9, an N-acetylserotonin methyltransferase gene, increases melatonin biosynthesis and improves water-use efficiency in transgenic apple. Tree physiology, 42, 1114–1126.